



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Preventing Fractures and Falls: A Limited Role for Calcium and Vitamin D Supplements?

Bischoff-Ferrari, Heike A ; Bhasin, Shalender ; Manson, JoAnn E

DOI: <https://doi.org/10.1001/jama.2018.4023>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-159504>

Journal Article

Published Version

Originally published at:

Bischoff-Ferrari, Heike A; Bhasin, Shalender; Manson, JoAnn E (2018). Preventing Fractures and Falls: A Limited Role for Calcium and Vitamin D Supplements? JAMA : the Journal of the American Medical Association, 319(15):1552-1553.

DOI: <https://doi.org/10.1001/jama.2018.4023>

Preventing Fractures and Falls

A Limited Role for Calcium and Vitamin D Supplements?

Heike A. Bischoff-Ferrari, MD, DrPH; Shalender Bhasin, MBBS; JoAnn E. Manson, MD, DrPH

Vitamin D and calcium are key nutrients to support bone development in children and young adults and to maintain muscle strength and neuromuscular coordination throughout the life span. Vitamin D deficiency is a well-defined risk factor for falls and hip fractures among older adults. It is unclear if supplementation with these nutrients is effective for the primary prevention of fractures among healthy community-dwelling adults.

In this issue of *JAMA*, the US Preventive Services Task Force (USPSTF) presents its recommendation statement on vitamin D and calcium supplementation to prevent fractures.¹ These recommendations were based on a careful and comprehensive



Related articles [pages 1592](#)
and [1600](#)

review of the evidence to determine whether supplements containing vitamin D or calcium individually or in combination contribute to the primary prevention of fractures among community-dwelling, asymptomatic adults without a diagnosis of osteoporosis or vitamin D deficiency.² In a simultaneously published report, the USPSTF also presents its recommendation statement on interventions to prevent falls in older adults,³ based on a comprehensive review of the evidence regarding strategies for fall prevention among community-dwelling adults 65 years or older, extending from vitamin D to exercise and multifactorial interventions.⁴ Providing both guidelines in tandem is valuable, as the complementary articles reinforce the importance of fall prevention in reducing the risk of fractures (and other injuries) among older adults, who sustain 75% of all osteoporotic fractures.⁵

Acknowledging the limited data available for primary prevention, the task force concluded “the current evidence is insufficient to assess the balance of the benefits and harms of vitamin D and calcium supplementation, alone or combined, for the primary prevention of fractures in community-dwelling, asymptomatic men and premenopausal women (I statement).”¹ It also recommended “against daily supplementation with 400 IU or less of vitamin D and 1000 mg or less of calcium for the primary prevention of fractures in community-dwelling, postmenopausal women (D recommendation).”¹ For daily supplementation with vitamin D doses greater than 400 IU and calcium doses greater than 1000 mg, the USPSTF concluded that “the current evidence is insufficient to assess the balance of the benefits and harms” for primary prevention of fractures in community-dwelling, postmenopausal women (I statement),¹ based on 2 large trials with inconsistent results.^{6,7} For the higher dose of vitamin D, the USPSTF identified 1 large trial that tested 100 000 IU of vitamin D every 4 months and demonstrated a benefit on fracture risk⁶ and 1 large trial that tested 100 000 IU of vitamin D monthly and did not find a benefit on fracture risk.⁷

With regard to possible harms of supplementation with vitamin D and calcium, the USPSTF found sufficient evidence that supplementation with calcium and vitamin D increases the risk of kidney stones, but identified no clear evidence for an increased risk of cardiovascular disease with vitamin D alone. The relationship between the dose of both nutrients and the risk of incident kidney stones could not be evaluated; in both of the relevant trials,^{8,9} however, the combined calcium intake from study medication and personal intake outside of the study was relatively high, often more than 2000 mg per day.

In the companion recommendation statement,³ the USPSTF updated and revised its earlier favorable assessment of vitamin D supplementation for fall prevention in 2012¹⁰ to a current recommendation “against vitamin D supplementation to prevent falls in community-dwelling adults 65 years or older (D recommendation).” The latter conclusion was based on inconsistent findings from 5 trials and likely influenced by evidence from a single trial that tested a large bolus dose of 500 000 IU of vitamin D annually¹¹ and reported increased risks of both falls and fractures with this treatment among community-dwelling adults at increased risk of falling. As alternative strategies, the USPSTF panel recommended, with moderate evidence, “exercise interventions to prevent falls in community-dwelling adults 65 years or older who are at increased risk for falls (B recommendation).”³ The exercise interventions evaluated were heterogeneous. Some exploratory analyses, which should be interpreted with caution, suggested that interventions that included group-based exercise, those with multiple components, and interventions including a strength or resistance component may be more effective. Common components of effective exercise interventions included training related to gait, balance, functional status, strength, flexibility, and endurance. The most common frequency of exercise interventions was 3 sessions per week.

The USPSTF recommendation, with increased emphasis on exercise, warrants adoption and should prove helpful, especially because exercise interventions reduce injurious falls. Moreover, the emphasis on physical activity should improve general health by reducing risks of other chronic diseases of aging, including coronary heart disease, stroke, type 2 diabetes, vascular dementia, and even cancer, as well as reducing the burden of mobility disability.^{12,13} A further recommendation, albeit with a lower level of evidence, concerned the implementation of multifactorial interventions. These interventions include an initial comprehensive assessment of modifiable risk factors for falls, such as the evaluation of balance, gait, vision, medication use, postural blood pressure, cognitive and psychological health, and

environmental factors. An individualized and tailored intervention is then developed, targeted to identified deficits. Multifactorial interventions may include components such as correction of vision or hearing deficits, nutritional therapy, medication management, home or environmental modifications, psychosocial interventions, and cognitive training, as well as exercise. Such multifactorial interventions were found to reduce the number of falls but not the number of falls with injuries.

Notably, the USPSTF recommendations apply only to community-dwelling and asymptomatic adults not known to have osteoporosis or vitamin D deficiency or to be at high risk of fracture. Thus, for patients at increased risk for osteoporosis, those with vitamin D deficiency, or both, it remains reasonable to consider vitamin D supplementation (800-1000 IU/d or more), consistent with recommendations of other professional societies such as the US Endocrine Society¹⁴ and National Osteoporosis Foundation.¹⁵ The literature is also supportive of a benefit of vitamin D supplementation in vulnerable populations, such as older adults living in institutions, other older adults at high risk of fracture, and those with vitamin D deficiency.^{16,17} The presence of major risk factors, such as a previous history of falls, age 75 years or older, and problems with mobility, gait, or balance, can help identify older adults at increased risk for fall injuries.^{3,18} A brief assessment of physical function using a Timed Up and Go test or a Short Physical

Performance Battery can also be helpful for identifying high-risk patients.^{3,18}

The USPSTF concluded that more evidence is needed to determine whether higher doses of vitamin D supplementation may be beneficial for primary fracture or fall prevention. To that regard, there will be a substantial increase in the available data on vitamin D supplementation among community-dwelling adults over the next year from 2 large double-blind randomized clinical trials (Vitamin D and Omega-3 Trial [VITAL; [NCT01169259](#)] in the United States and DO-HEALTH [[NCT01745263](#)] in Europe), both testing 2000 IU daily vitamin D₃ vs placebo. In addition, 2 other large ongoing clinical trials are expected to present their findings within 2 years (FIND in Finland, testing 1600 IU and 3200 IU vitamin D daily vs placebo [[NCT01463813](#)], and D-Health in Australia, testing 60 000 IU of vitamin D monthly vs placebo [[ACTRN12613000743763](#)]). The European DO-HEALTH trial, being conducted among healthy adults 70 years or older, is also testing a home-based exercise and strength program for the prevention of both falls and fractures. Last, 2 large cluster randomized pragmatic trials, Strategies to Reduce Injuries and Develop Confidence in Elders (STRIDE; [NCT02475850](#)) and the Prevention of Fall Injuries Trial (PreFIT; [ISRCTN71002650](#)), are evaluating the clinical effectiveness of multifactorial strategies in preventing fall injuries. Thus, clinicians will soon have more data to guide their treatment decisions in these important areas.

ARTICLE INFORMATION

Author Affiliations: Department of Geriatrics and Aging Research, University Hospital Zurich and Waid City Hospital, University of Zurich, Zurich, Switzerland (Bischoff-Ferrari); Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts (Bhasin, Manson); Research Program in Men's Health: Aging and Metabolism, and the Boston Claude D. Pepper Older Americans Independence Center, Boston, Massachusetts (Bhasin); Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Manson).

Corresponding Author: JoAnn E. Manson, MD, DrPH, Division of Preventive Medicine, Brigham and Women's Hospital, Harvard Medical School, 900 Commonwealth Ave E, Boston, MA 02215 (jmanson@rics.bwh.harvard.edu).

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Dr Bhasin reported receiving grant funding from the National Institute on Aging, the National Institute of Nursing Research, AbbVie, MIB LLC, AliveGen, Abbott, and Transition Therapeutics; receiving personal fees from Novartis; holding equity interest in FPT LLC; and serving as chair of the American Board of Internal Medicine examination writing committee for endocrinology. Dr Manson reported receiving grant funding from the National Institutes of Health (NIH) and serving as a principal investigator of the VITamin D and Omega-3 Trial (VITAL), which is funded by the NIH. No other disclosures were reported.

REFERENCES

1. US Preventive Services Task Force. Vitamin D, calcium, or combined supplementation for the primary prevention of fractures in community-

dwelling adults: US Preventive Services Task Force recommendation statement [published online April 17, 2018]. *JAMA*. doi:10.1001/jama.2018.3185

2. Kahwati LC, Weber RP, Pan H, et al. Vitamin D, calcium, or combined supplementation for the primary prevention of fractures in community-dwelling adults: evidence report and systematic review for the US Preventive Services Task Force [published online April 17, 2018]. *JAMA*. doi:10.1001/jama.2017.21640

3. US Preventive Services Task Force. Interventions to prevent falls in community-dwelling older adults: US Preventive Services Task Force recommendation statement [published online April 17, 2011]. *JAMA*. doi:10.1001/jama.2018.3097

4. Guirguis-Blake JM, Michael YL, Perdue LA, et al. Interventions to prevent falls in community-dwelling older adults: updated evidence report and systematic review for the US Preventive Services Task Force [published online April 17, 2018]. *JAMA*. doi:10.1001/jama.2017.21962

5. Melton LJ III, Crowley CS, O'Fallon WM. Fracture incidence in Olmsted County, Minnesota. *Osteoporos Int*. 1999;9(1):29-37.

6. Trivedi DP, Doll R, Khaw KT. Effect of four monthly oral vitamin D₃ (cholecalciferol) supplementation on fractures and mortality in men and women living in the community. *BMJ*. 2003;326(7387):469.

7. Khaw KT, Stewart AW, Waayer D, et al. Effect of monthly high-dose vitamin D supplementation on falls and non-vertebral fractures. *Lancet Diabetes Endocrinol*. 2017;5(6):438-447.

8. Jackson RD, LaCroix AZ, Gass M, et al; Women's Health Initiative Investigators. Calcium plus vitamin D supplementation and the risk of fractures. *N Engl J Med*. 2006;354(7):669-683.

9. Lappe J, Watson P, Travers-Gustafson D, et al. Effect of vitamin D and calcium supplementation on cancer incidence in older women. *JAMA*. 2017;317(12):1234-1243.

10. Moyer VA; U.S. Preventive Services Task Force. Prevention of falls in community-dwelling older adults. *Ann Intern Med*. 2012;157(3):197-204.

11. Sanders KM, Stuart AL, Williamson EJ, et al. Annual high-dose oral vitamin D and falls and fractures in older women. *JAMA*. 2010;303(18):1815-1822.

12. US Department of Health and Human Services. 2008 Physical Activity Guidelines for Americans. Washington, DC: US Dept of Health and Human Services; 2008.

13. Pahor M, Guralnik JM, Ambrosius WT, et al; LIFE Study Investigators. Effect of structured physical activity on prevention of major mobility disability in older adults. *JAMA*. 2014;311(23):2387-2396.

14. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al; Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency. *J Clin Endocrinol Metab*. 2011;96(7):1911-1930.

15. Cosman F, de Beur SJ, LeBoff MS, et al; National Osteoporosis Foundation. Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int*. 2014;25(10):2359-2381.

16. Bischoff-Ferrari HA, Willett WC, Orav EJ, et al. A pooled analysis of vitamin D dose requirements for fracture prevention. *N Engl J Med*. 2012;367(1):40-49.

17. Weaver CM, Alexander DD, Boushey CJ, et al. Calcium plus vitamin D supplementation and risk of fractures. *Osteoporos Int*. 2016;27(1):367-376.

18. Bhasin S, Gill TM, Reuben DB, et al. Strategies to Reduce Injuries and Develop Confidence in Elders (STRIDE). *J Gerontol A Biol Sci Med Sci*. 2017. doi:10.1093/gerona/glx190